



This is an official  
**CDC Health Advisory**

Distributed via Health Alert Network  
January 17, 2013, 10:45AM  
10275-CHA-01-17-2013-FLU

## **Notice to Clinicians: Summary of CDC Recommendations for Influenza Antiviral Medications**

CDC continues to recommend antiviral medications for treatment of seasonal influenza and annual vaccination as the best tools for prevention<sup>1</sup>.

Evidence from past influenza seasons and the 2009 H1N1 pandemic has shown that treatment with antiviral medications can have clinical and public health benefit in reducing severe outcomes of influenza when initiated as soon as possible after illness onset.

Clinical trials and observational data show that early antiviral treatment may do the following:

- shorten the duration of fever and illness symptoms
- reduce the risk of complications from influenza (e.g., otitis media in young children, pneumonia, respiratory failure) and death
- shorten the duration of hospitalization

Below is a summary of CDC's influenza antiviral recommendations.

### **Summary of CDC recommendations for influenza antiviral medications for the 2012-2013 season:**

Clinical benefit is greatest when antiviral treatment is administered early. When indicated, antiviral treatment should be started as soon as possible after illness onset, ideally within 48 hours of symptom onset. However, antiviral treatment might still be beneficial in patients with severe, complicated, or progressive illness and in hospitalized patients when started after 48 hours of illness onset, as indicated by observational studies.

Antiviral treatment is recommended as early as possible for any patient with confirmed or suspected influenza who

- is hospitalized;
- has severe, complicated, or progressive illness; or
- is at higher risk for influenza complications. This list includes:
  - children aged younger than 2 years;<sup>2</sup>
  - adults aged 65 years and older;
  - persons with chronic pulmonary (including asthma), cardiovascular (except hypertension alone), renal, hepatic, hematological (including sickle cell disease), metabolic disorders (including diabetes mellitus), or neurologic and neurodevelopment conditions (including disorders of the brain, spinal cord, peripheral nerve, and muscle such as cerebral palsy, epilepsy [seizure disorders], stroke, intellectual disability [mental retardation], moderate to severe developmental delay, muscular dystrophy, or spinal cord injury);

- persons with immunosuppression, including that caused by medications or by HIV infection;
- women who are pregnant or postpartum (within 2 weeks after delivery);
- persons aged younger than 19 years who are receiving long-term aspirin therapy;
- American Indians/Alaska Natives;
- persons who are morbidly obese (i.e., body-mass index is equal to or greater than 40); and
- residents of nursing homes and other chronic-care facilities.

Clinical judgment, on the basis of the patient's disease severity and progression, age, underlying medical conditions, likelihood of influenza, and time since onset of symptoms, is important when making antiviral treatment decisions for high-risk outpatients.

Decisions about starting antiviral treatment should not wait for laboratory confirmation of influenza.

While influenza vaccination is the first and best way to prevent influenza, a history of influenza vaccination does not rule out the possibility of influenza virus infection in an ill patient with clinical signs and symptoms compatible with influenza.

Antiviral treatment also can be considered for any previously healthy, symptomatic outpatient not at high risk with confirmed or suspected influenza on the basis of clinical judgment, if treatment can be initiated within 48 hours of illness onset.

#### **For more information:**

*A full summary of clinical recommendations that includes the sections listed below is available at*

<http://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>:

- [Antiviral Medications Recommended for Treatment and Chemoprophylaxis of Influenza](#)
- [Summary of Influenza Antiviral Treatment Recommendations](#)
- [Diagnostic Testing for Influenza](#)
- [Recommended Dosage and Duration of Treatment or Chemoprophylaxis for Influenza Antiviral Medications](#)
- [Chemoprophylaxis](#)
- [Adverse Events](#)
- [Footnotes](#)
- [Selected References](#)

*CDC communication to health care providers:*

- CDC Information for Pharmacists <http://www.cdc.gov/flu/professionals/2012-2013-guidance-pharmacists.htm>
- CDC Influenza Update for Pediatricians <http://www.cdc.gov/flu/professionals/2012-2013-guidance-pediatricians.htm>

*FDA Announcement Addressing Intermittent Shortages of Oseltamivir Phosphate (Tamiflu®) for Oral Suspension (6mg/mL 60 mL):*

On January 10, 2013, the U.S. Food and Drug Administration (FDA) released information indicating there may currently be intermittent shortages of [Oseltamivir Phosphate \(Tamiflu®\) for Oral Suspension \(6mg/mL 60 mL\)](#) - used to treat influenza in children - due to increased demand for the drug. This is the pediatric suspension (liquid). Instructions for pharmacists on how [to compound an oral suspension from Tamiflu® 75 mg \(adult\) capsules](#) are available at [http://www.tamiflu.com/hcp/resources/hcp\\_resources\\_pharmacists.jsp](http://www.tamiflu.com/hcp/resources/hcp_resources_pharmacists.jsp). These instructions provide for an

alternative oral suspension when commercially manufactured oral suspension formulation is not readily available.

In some cases, clinicians may consider substituting a 30 or 45 mg capsule for children (if dose is appropriate) rather than suspension, particularly if there are spot shortages of suspension. These capsules may be opened and mixed with a sweet liquid, such as regular or sugar-free chocolate syrup, if oral suspension is not available. Instructions to share with caregivers are available here:

[www.cdc.gov/flu/antivirals/mixing\\_tamiflu\\_qa.htm](http://www.cdc.gov/flu/antivirals/mixing_tamiflu_qa.htm) .

---

<sup>i</sup> Findings from early data indicate the overall effectiveness of the 2012-2013 seasonal influenza vaccine is 62%. The data are published in “[Early Estimates of Seasonal Influenza Vaccine Effectiveness — United States, January 2013](#),” in the January 11, 2013, Morbidity and Mortality Weekly Report. This estimate is within the range of what is expected during seasons when most circulating influenza viruses characterized by CDC are like the viruses included in the vaccine.

At this time, people seeking vaccination may need to call more than one provider to locate vaccine. The [flu vaccine locator](http://flushot.healthmap.org/) (<http://flushot.healthmap.org/>) may be helpful.

<sup>ii</sup> On December 21, 2012, the U.S. Food and Drug Administration (FDA) approved the antiviral medication oseltamivir (trade name Tamiflu®) for the treatment of influenza in people aged 2 weeks and older. An FDA press release related to this announcement is available at <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm333205.htm>

## DHEC contact information for reportable diseases and reporting requirements

Reporting of influenza is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2013 List of Reportable Conditions available at:

<http://www.scdhec.gov/health/disease/reportables.htm>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

### Regional Public Health Offices – 2013

**Mail or call reports to the Epidemiology Office in each Public Health Region.**

#### Region 1

##### **Anderson, Oconee**

220 McGee Road  
Anderson, SC 29625  
Phone: (864) 260-4358  
Fax: (864) 260-5623  
Nights / Weekends: 1-866-298-4442

##### **Abbeville, Edgefield, Greenwood, Laurens, McCormick, Saluda**

1736 S. Main Street  
Greenwood, SC 29646  
Phone: 1-888-218-5475  
Fax: (864) 942-3690  
Nights / Weekends: 1-800-420-1915

#### Region 2

##### **Greenville, Pickens**

PO Box 2507  
200 University Ridge  
Greenville, SC 29602-2507  
Phone: (864) 282-4139  
Fax: (864) 282-4373  
Nights / Weekends: (864) 809-3825

##### **Cherokee, Spartanburg, Union**

PO Box 4217  
151 E. Wood Street  
Spartanburg, SC 29305-4217  
Phone: (864) 282-4139  
Fax: (864) 282-4373  
Nights / Weekends: (864) 809-3825

#### Region 3

##### **Chester, Lancaster, York**

PO Box 817  
1833 Pageland Highway  
Lancaster, SC 29720  
Phone: (803) 286-9948  
Fax: (803) 286-5418  
Nights / Weekends: 1-866-867-3886

#### Region 3 (continued)

##### **Fairfield, Lexington, Newberry, Richland**

2000 Hampton Street  
Columbia, SC 29204  
Phone: (803) 576-2749  
Fax: (803) 576-2993  
Nights / Weekends: 1-888-554-9915

#### Region 4

##### **Chesterfield, Darlington, Dillon, Florence, Marlboro, Marion**

145 E. Cheves Street  
Florence, SC 29506  
Phone: (843) 661-4830  
Fax: (843) 661-4859  
Nights / Weekends: (843) 601-7051

##### **Clarendon, Kershaw, Lee, Sumter**

PO Box 1628  
105 North Magnolia Street  
Sumter, SC 29150  
Phone: (803) 773-5511  
Fax: (803) 775-9941  
Nights/Weekends: 803-458-1847

#### Region 5

##### **Bamberg, Calhoun, Orangeburg**

PO Box 1126  
1550 Carolina Avenue  
Orangeburg, SC 29116  
Phone: (803) 533-7199  
Fax: (803) 533-7134  
Nights / Weekends: (803) 535-9265

##### **Aiken, Allendale, Barnwell**

222 Beaufort Street, NE  
Aiken, SC 29801  
Phone: (803) 642-1618  
Fax: (803) 643-8386  
Nights / Weekends: (803) 645-8167

#### Region 6

##### **Georgetown, Horry, Williamsburg**

1931 Industrial Park Road  
Conway, SC 29526-5482  
Phone: (843) 915-8804  
Fax: (843) 365-0085  
Nights/Weekends: (843) 340-4754

#### Region 7

##### **Berkeley, Charleston, Dorchester**

4050 Bridge View Drive, Suite 600  
N. Charleston, SC 29405  
Phone: (843) 953-0047  
Fax: (843) 953-0051  
Nights / Weekends: (843) 219-8470

#### Region 8

##### **Beaufort, Colleton, Hampton, Jasper**

219 S. Lemacks Street  
Walterboro, SC 29488  
Phone: (843) 441-1091  
Fax: (843) 549-6845  
Nights / Weekends: 1-843-441-1091

#### DHEC Bureau of Disease Control

##### **Division of Acute Disease Epidemiology**

1751 Calhoun Street  
Box 101106  
Columbia, SC 29211  
Phone: (803) 898-0861  
Fax: (803) 898-0897  
Nights / Weekends: 1-888-847-0902



PROMOTE PROTECT PROSPER  
South Carolina Department of Health  
and Environmental Control

[www.scdhec.gov](http://www.scdhec.gov)

Categories of Health Alert messages:

<b>Health Alert</b>	Conveys the highest level of importance; warrants immediate action or attention.
<b>Health Advisory</b>	Provides important information for a specific incident or situation; may not require immediate action.
<b>Health Update</b>	Provides updated information regarding an incident or situation; unlikely to require immediate action.